

Anti-Human BSX, monoclonal

Alternate Names: Brain-specific homeobox protein homolog

Cat. No. 13-067

FOR RESEARCH USE ONLY

NOT FOR USE IN HUMANS



PRODUCT DESCRIPTION

UniProt Summary

UniProt

Primary Accession: [Q3C1V8](#)

DNA binding protein that function as transcriptional activator. Is essential for normal postnatal growth and nursing. Is an essential factor for neuronal neuropeptide Y and agouti-related peptide function and locomotory behavior in the control of energy balance (By similarity). Nucleus (By similarity). Belongs to the distal-less homeobox family. Contains 1 homeobox DNA-binding domain. Name=Protein Spotlight; Note=Of fidgets and food - Issue 85 of August 2007; URL="http://web.expasy.org/spotlight/back_issues/0;

Physical Characteristics

Quantity: 1 ml (culture supernatant) or 100ug at 1mg/ml (purified)

Format: culture supernatant or purified material

Host/Isotype: mouse IgG2b

Clonality: monoclonal; ID R329.1.5H1

Formulation: culture supernatant contains 0.02% NaN₃. Purified material contains 30% glycerol, PBS and 0.02% NaN₃

Specificity: monospecific for human BSX ; see "Microarray Analysis" below

Reactivity: human; not tested for cross reactivity in other species

Stability/Storage: 12 months long term: -20°C; short term: 4°C; avoid freeze-thaw cycles; aliquot as required

Handling Notes: small volumes of antibody may occasionally become entrapped in the seal of the product vial during shipment and storage; briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the container cap

Tested Research Applications

- Western Immunoblotting - PASS by SOP
- Immunoprecipitation - PASS by SOP

Antibody tested as purified IgG. Optimal dilution to be determined by user.

Quality Assurance

Notes:

1. Please refer to the SOP manual ([click here](#)) to for detailed explanation of the experimental setup and thresholds used to evaluate the results shown below.
2. Predicted MW of the target protein expressed in the western immunoblotting (WB) and immprecipitation (IP) experiment is **25.74kDa** and the three fusion tags (venus, 3xFLAG, V5) adds another **38.7kDa** to this protein. Therefore the protein is expected to migrate **~64.44kDa** in a denatured SDS-PAGE gel.
3. See results below for any applicable cautionary notes.

Guanajibo Research and Innovation Park
4005 St B Road 114 Km 1.3
Mayaguez, PR 00682

T 787.806-4100 Ext 233
F 787.806-4006
www.cdi-lab.com

© CDI Laboratories 2017.
For research use only. Not for use in humans or for diagnostics.
Lot-specific COA version tracker: v1.0.0



Anti-Human BSX, monoclonal

Alternate Names: Brain-specific homeobox protein homolog

Cat. No. 13-067


FOR RESEARCH USE ONLY

NOT FOR USE IN HUMANS



PRODUCT DESCRIPTION

Quality Assurance (HuProt™ Array)



About Z and S Scores: The z-score represents the strength of a signal that a mAb (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. To be considered a definite binding event between a mAb and a protein target, the z-score must have a value of at least 2.5 (SDs above the mean). The s-score represents the relative target specificity of a mAb, and is the difference (also in units of SDs) between the z-score a mAb generates upon binding to one protein and the next highest z-score that the mAb generates against another protein. For example, if a mAb binds to protein X with an z-score of 43 and to protein Y with an z-score of 14, then the s-score for the binding of that mAb to protein X is equal to 29.

Guanajibo Research and Innovation Park
4005 St B Road 114 Km 1.3
Mayaguez, PR 00682

T 787.806-4100 Ext 233
F 787.806-4006
www.cdi-lab.com

© CDI Laboratories 2017.
For research use only. Not for use in humans or for diagnostics.
Lot-specific COA version tracker: v1.0.0



Anti-Human BSX, monoclonal

Alternate Names: Brain-specific homeobox protein homolog

Cat. No. 13-067

FOR RESEARCH USE ONLY

NOT FOR USE IN HUMANS




NEXTGEN PROTEOMICS

PRODUCT DESCRIPTION

Quality Assurance (continued)

IMMUNOPRECIPITATION - Predicted MW 64.44kDa WESTERN BLOT - Predicted MW 64.44kDa



Tet-ON HeLa cells were transfected with construct encoding BSX (NM_001098169.1) with an N-terminal fusion of FLAG, YFP (Venus) and V5 tags under a tet-inducible promoter. These cells were stimulated with 0, 50 or 1000 ng/ml doxycycline. Immunoprecipitation (IP) was carried out using 5 μ g of either IgG, CDI mAb Anti-BSX (cloneID# R329.1.5H1) or 1 μ g of FLAG-M2. Immunoblotting was performed using rabbit Anti-FLAG (1:1000, Cell Signaling #2368).

Tet-ON HeLa cells were transfected with construct encoding BSX (NM_001098169.1) with an N-terminal fusion of FLAG, YFP (Venus) and V5 tags under a tet-inducible promoter. These cells were stimulated with 0, 50 or 1000 ng/ml doxycycline. Immunoprecipitation (IP) was carried out using 5 μ g of either IgG, CDI mAb Anti-BSX (cloneID# R329.1.5H1) or 1 μ g of FLAG-M2. Immunoblotting was performed using 0.2 μ g/ml CDI mouse mAb Anti-BSX (cloneID# R329.1.5H1). HC=Heavy chain.

Cautionary notes for IP and WB experiments:

1. No or low signal seen in Flag IP lane

Guanajibo Research and Innovation Park
4005 St B Road 114 Km 1.3
Mayaguez, PR 00682

T 787.806-4100 Ext 233
F 787.806-4006
www.cdi-lab.com

© CDI Laboratories 2017.
For research use only. Not for use in humans or for diagnostics.
Lot-specific COA version tracker: v1.0.0



NEXTGEN PROTEOMICS